RYPOS

Understanding Clean Air: Defining Purity and Identifying Pollutants

Ian Sines
Research Manager
Rypos

Introduction

Clean air is something we rarely think about, until it's gone. Whether indoors or out, the air we breathe directly affects our health, comfort, and productivity. Yet defining what makes air “clean” isn’t always straightforward. While regulatory agencies set standards for specific pollutants, real-world air quality is shaped by a broader mix of particles, gases, and odors. This paper explores the full scope of what clean air means today, how contamination occurs, and the technologies available to protect and improve the air around us.

What Is Clean Air?

Clean air is fundamental to public health, environmental protection, and overall quality of life. The U.S. Environmental Protection Agency (EPA) defines clean air as air containing pollutant concentrations below the thresholds established by the National Ambient Air Quality Standards (NAAQS). These standards target six key pollutants known to pose serious health and environmental risks [1]:

- Particulate Matter (PM)
- Nitrogen Dioxide (NO_2)
- Ozone (O_3)
- Sulfur Dioxide (SO_2)
- Carbon Monoxide (CO)
- Lead (Pb)

The EPA further stipulates that clean air must be safe to breathe for everyone—including children, the elderly, and those with pre-existing respiratory conditions. Beyond human health, clean air must also protect the environment from harmful effects such as acid rain, smog, and damage to vegetation.

However, practical definitions of clean air often extend beyond NAAQS criteria. Many airborne contaminants, while not regulated by federal standards, can still degrade air quality. Odors from sources such as sewage, mold, sweat, garbage, or decomposing organic matter (e.g., trimethylamine from rotting fish) can render air unbreathable or offensive.

Airborne pathogens, such as influenza viruses contained in respiratory droplets, are also a form of contamination with serious health implications.[2] In short, many types of air contamination exist beyond EPA definitions and require comprehensive mitigation strategies.

Understanding Air Contaminants

Air contamination can be broadly categorized into two main types:

1. **PM Contamination** [3]

PM consists of a mix of solid particles and liquid droplets suspended in air. These particles vary in size and composition and originate from both anthropogenic (human-made) and natural sources.

- **Anthropogenic sources**: Diesel engine soot, construction dust, industrial emissions.
- **Natural sources**: Pollen, fungal spores, wildfire smoke, volcanic ash.

PM is particularly dangerous due to its ability to penetrate the respiratory system. Fine particles ($PM_{2.5}$) can lodge deep in the lungs or even enter the bloodstream, contributing to conditions such as asthma, heart disease, and lung cancer.

2. **MC Control** [4]

Molecular contaminants (MC) are airborne gases or vapors that are fully dissolved in the air. They can be toxic, corrosive, flammable, or odorous. Common examples include:

- Ammonia (NH_3)
- Hydrogen Sulfide (H_2S)
- Volatile Organic Compounds (VOCs)
- Formaldehyde
- Alcohol and Ketones

To illustrate: a household cat can contribute to both PM and MC contamination. Dander (PM) may trigger allergic reactions, while compounds emitted from a litterbox (MC), such as ammonia and sulfur-based gases, can quickly degrade indoor air quality and create an environment that is unpleasant and degrades indoor air quality.

Mitigating Air Contamination

The mitigation of air pollution depends on understanding the characteristics of the contaminants and applying appropriate control technologies. Effective solutions typically fall into separate categories for PM and MC.

Particulate Matter Control

Several technologies exist to mitigate PM, depending on the size and nature of the particles, the volume of air being treated, and the application environment.

- **Mechanical Filtration**: Devices like HEPA (High Efficiency Particulate Air) filters use dense media to trap PM. These are widely used in indoor environments and medical settings.
- **Electrostatic Precipitation**: This method applies an electric field to ionize particles and collect them on charged plates. It is commonly used in industrial emissions control.

Molecular Contamination Control

Unlike PM, molecular pollutants require different mitigation techniques:

- **Adsorption Filtration**: High surface area materials such as activated carbon, zeolites, or silicas absorb gaseous molecules. These filters act like sponges and are particularly effective against VOCs and odors. In some cases, these systems can be regenerated and reused, such as in rotary concentrator systems.
- **Thermal Oxidation and Catalysis**: Catalytic converters are a prime example of this strategy, where high temperatures and noble metal catalysts (e.g., platinum, palladium)

oxidize harmful gases into less toxic compounds. For instance, CO is converted to carbon dioxide (CO_2), a significantly less hazardous emission. These systems, when operated correctly, can offer long service lives with minimal maintenance.

- **Biological Filtration**: Emerging as a novel approach, biofilters use microorganisms to break down both PM and MC. While potentially sustainable and long-lasting, these systems are sensitive to environmental extremes and may fail in the presence of certain chemicals or temperatures.

Emissions Reduction Technologies: Advancing Clean Air Across Applications

Reducing emissions from combustion sources—especially diesel engines—requires a multifaceted approach, often involving a combination of filtration, chemical conversion, and system design optimization. Several established and emerging technologies are used to target both PM and MC in stationary and mobile applications.

Particulate Matter Control Technologies

- **Passive Diesel Particulate Filters (DPFs)**: These filters capture soot and rely on high exhaust temperatures to regenerate (burn off) accumulated PM. They are most effective in applications with consistent load profiles and sufficient exhaust heat.
- **Active Diesel Particulate Filters**: These systems include an external heat source or electrical element to initiate regeneration, making them suitable for variable-load or low-temperature operations such as standby generators or cold starts.[5]
- **High-Efficiency Filters with Regeneration Cycles**: Some advanced DPFs incorporate real-time monitoring and automatic regeneration to maintain low back pressure and maximize uptime. These are used in critical infrastructure applications requiring reliability and emissions compliance.

Molecular Contamination and Gaseous Emission Controls

- **Selective Catalytic Reduction (SCR)**: This technology reduces nitrogen oxides (NO_x) by injecting a urea-based solution into the exhaust stream. In the presence of a catalyst, NO_x is converted to nitrogen and water vapor. SCR is commonly used in conjunction with DPFs in Tier 4 systems.[6]
- **Oxidation Catalysts (DOCs)**: Diesel oxidation catalysts help reduce CO, unburned hydrocarbons (HC), and certain soluble organic fractions of PM. These catalysts use precious metals to facilitate chemical conversion at relatively low temperatures.
- **Adsorption Systems**: Activated carbon and other porous media are used in some systems to control odor and VOCs not typically addressed by traditional combustion aftertreatment technologies.[7]
- **Hybrid and Integrated Systems**: Many emissions reduction solutions combine multiple technologies—such as DPF + SCR or DOC + DPF systems—to meet stringent regulatory limits while optimizing performance, maintenance, and cost.

These technologies continue to evolve as regulatory pressures tighten and new use cases emerge, including microgrids, renewable energy backup, and ultra-low emission zones.

Selecting the right emissions control solution depends on application-specific factors such as engine size, operating cycle, load profile, and environmental requirements.

Clean Air is a Public Health Imperative

The definition of clean air extends beyond regulatory thresholds—it includes eliminating contaminants that impact health, comfort, and environmental quality. Whether dealing with particulates, gases, odors, or biological threats, the right mitigation technologies can make the difference between clean and polluted air.

Understanding the sources and behavior of airborne contaminants allows engineers, regulators, and facility operators to apply targeted control solutions—ranging from filtration to catalytic conversion to biological treatment. As the demand for cleaner, more resilient systems grows, so too must our commitment to innovation in air quality and emissions control technologies.

References

[1] U.S. Environmental Protection Agency, “Naaqs Table”, from https://www.epa.gov/criteria-air-pollutants/naaqs-table

[2] La Rosa, Giuseppina; Fratini, Marta; Della Libera, Simonetta; Iaconelli, Marcello; Muscillo, Michele (2013-06-01). "Viral infections acquired indoors through airborne, droplet or contact transmission". *Annali dell'Istituto Superiore di Sanità*. **49** (2): 124–132. doi:10.4415/ANN_13_02_03. ISSN 0021-2571. PMID 23771256

[3] *Seinfeld JH, Pandis S (1998). Atmospheric Chemistry and Physics: From Air Pollution to Climate Change (2nd ed.). Hoboken, New Jersey: John Wiley & Sons. p. 97. ISBN 978-0-471-17816-3.*

[4] Davies, Helen L.; O'Leary, Catherine; Dillon, Terry; Shaw, David R.; Shaw, Marvin; Mehra, Archit; Phillips, Gavin; Carslaw, Nicola (August 14, 2023). "A measurement and modelling investigation of the indoor air chemistry following cooking activities". *Environmental Science: Processes & Impacts*. **25** (9): 1532–1548. doi:10.1039/D3EM00167A. ISSN 2050-7887. PMID 37609942.

[5] Rypos, “Diesel Particulate Filter (DPF) Exhaust Systems”, from https://www.rypos.com/diesel-particulate-filter/

[6] Han, Lupeng; Cai, Sixiang; Gao, Min; Hasegawa, Jun-ya; Wang, Penglu; Zhang, Jianping; Shi, Liyi; Zhang, Dengsong (2019), from "Selective Catalytic Reduction of NO_x with NH_3 by Using Novel Catalysts: State of the Art and Future Prospects". *Chemical Reviews*. **119** (19): 10916–10976.

[7] Chalmpes N, Ochonma P, Tantis I, Alsmaeil AW, Assafa TE, Tathacharya M, Srivastava M, Gadikota G, Bourlinos AB, Steriotis T, Giannelis EP (2024-12-10), from "Ultrahigh Surface Area Nanoporous Carbons Synthesized via Hypergolic and Activation Reactions for Enhanced CO_2 Capacity and Volumetric Energy Density". *ACS Nano*. **18** (49): 33491–33504. Bibcode:2024ACSNa..1833491C. doi:10.1021/acsnano.4c10531. ISSN 1936-0851. PMID 39576877.

2025 Rypos, Inc.
9/2025